

Is simultaneous y and ξ -scaling in the quasi-elastic region accidental?

Donal Day

Dept. of Physics, Univ. of Virginia, Charlottesville, VA, USA

Ingo Sick

Dept. für Physik und Astronomie, Univ. Basel, Basel, Switzerland

We study the y and ξ -scaling of the nuclear response at large momentum transfer in order to understand how scaling based on very different descriptions of the elementary interaction can occur simultaneously. We find that the approximate validity of ξ -scaling at low energy loss arises from the coincidental behavior of the quasielastic and deep inelastic cross sections.

I. INTRODUCTION

Scaling occurs in several fields where a probe scatters incoherently from a composite system: quasi-elastic scattering of electrons from nucleons bound in nuclei [1], deep inelastic scattering (DIS) of leptons from quarks bound in nucleons [2], scattering of thermal neutrons from individual atoms in liquids such as $L^4\text{He}$ [3, 4], and Compton scattering from atoms [5], to name a few. These inclusive scattering cross sections *a priori* are functions of two independent variables, the momentum transfer q and the energy transfer ν . The scattering is said to scale if the cross sections, after removal (by division) of the elementary probe-constituent cross section and a kinematic factor determined by the elastic scattering kinematics, become a function of a *single* variable $z = z(q, \nu)$ rather than q and ν separately [6]. The observation of scaling (and of scaling violations) provides a better understanding of both the reaction mechanism and the structure of the system in which the constituent is bound.

Quasi-elastic electron-nucleus scattering shows a scaling behavior once the inclusive scattering cross sections are divided by the elastic electron-nucleon cross section and a kinematic factor connecting the scaling variable and energy and momentum transfer. The corresponding scaling variable $y \equiv -q + (\nu^2 + 2m\nu)^{1/2}$ is obtained from energy/momentum conservation for elastic scattering from an initially off-shell nucleon, and the scaling function $F(y)$ represents the probability to find nucleons in the nucleus with the minimum momentum y allowed by the kinematics. The minimum momentum, y , is parallel to $\vec{q}$.

Scaling of the nucleon structure functions, MW_1 and νW_2 , in DIS is revealed after division of the inclusive cross section by the Mott cross section (describing the Q^2 dependence of point-like scattering). The scaling variable in DIS, $x = Q^2/2m\nu$, (Q is the four-momentum transfer) is the struck parton's fraction of the nucleon momentum in the infinite momentum frame. The scaling function νW_2 allows access to the quark distribution functions. The Nachtmann variable [7] $\xi = (q - \nu)/m$ (other equivalent expressions are customary in the literature) is often used to extend DIS scaling to lower momentum transfers.

In the kinematical regions dominated by quasi-elastic scattering and DIS, scaling is observed in y and x (or

ξ), respectively. Deviations from ideal scaling in quasi-elastic scattering are due to the effects of the final-state interaction (FSI) of the recoiling nucleon, and the importance of large removal energies E in the nuclear spectral function $S(k, E)$ [6]. For DIS, scaling violations mainly arise from the evolution of the structure functions with Q^2 and are well understood within perturbative QCD [8]. When neglecting FSI, νW_2 provides information on the quark distribution functions [9]. ξ , for a mass-zero constituent initially off-shell, is its momentum component parallel to $\vec{q}$, in units of the nucleon mass m .

ξ -scaling Recent work on ξ -scaling has been focused on exploring both resonance region and the DIS region. It has been found that the response averaged over the resonances defines the same scaling function as derived from DIS at very large momentum transfer and evolved to Q^2 appropriate for the resonances [10, 11]. This observation, first revealed observed some years ago, and referred to as “duality” [12], is only partly understood. The extension of such studies to very low energy loss, the region of the quasi-elastic peak in nuclei at $y < 0$ or $x > 1$, showed [13] that ξ -scaling apparently works even where the dynamics of probe-constituent scattering (quasi-elastic scattering from a nucleon) is very different from the one used in ξ -scaling.

The observation of ξ -scaling in the region of the low- ν side of the quasi-elastic peak certainly is puzzling. y -scaling in this region is dependent on the use of elastic electron-nucleon kinematics for the scattering process, which is quite different for DIS as the former involves a dependence on the finite nucleon mass (at the q 's of interest here). As importantly, the elementary probe-constituent cross section momentum dependences are quite dissimilar. In ξ -scaling the Mott cross section carries all the Q^2 dependence while for y -scaling the electron-nucleon cross sections depend on the nucleon elastic form factors which carry an additional strong Q^2 dependence.

Benhar and Liuti [14] already have noted this puzzle and suggested an explanation. For quasi-elastic scattering at moderately large q the FSI of the recoiling nucleon still plays an important role [15]. This FSI leads to a violation of y -scaling. Benhar and Liuti showed that, for the inclusive cross sections then available, the Q^2 dependences of the FSI and of the relationship between ξ and

y could combine such that there would be, accidentally, an approximate scaling in terms of ξ .

An experiment (E89-008) performed at Jefferson Lab provided a much more extensive set of data[16] on quasi-elastic electron-nucleus scattering. This data extends to much larger momentum transfers and, for a given y or ξ , covers a larger range in Q^2 . These data revealed, as expected, y -scaling on the low- ν side of the quasi-elastic peak. Surprisingly it also displayed an approximate scaling in terms of ξ which, at the largest q , extends to rather low values of ν . The authors of [16] pointed out that over this larger range of q the explanation for scaling in terms of ξ provided by Benhar and Liuti could no longer be valid; at large Q^2 the FSI decreases, while the elementary cross sections and the kinematics describing the reaction for ξ and y scaling remain disparate. The simultaneous scaling in y and ξ thus remains a puzzle, and it is not clear whether ξ scaling in the quasi-elastic region occurs due to some accidental convergence of factors or whether it has a deeper meaning.

In this Letter, we return to this question of “accidental” ξ -scaling to provide additional insight.

II. PIECES OF THE PUZZLE

An understanding of the simultaneous display of y and ξ scaling in the quasi-elastic region requires going beyond the question that was addressed by Benhar and Liuti [14]. The definitions of y and ξ provide part of the understanding as they lead to a relative shift of data points taken at different Q^2 's, but that can not explain the phenomenon. The y -scaling function, $F(y)$ is obtained by dividing the inclusive cross section by the cross section for incoherent electron-nucleon elastic scattering (summed over the number of protons and neutrons). In the case of the ξ -scaling function, νW_2 , only the Mott cross section is divided out. These two cross sections differ by the nucleon elastic form factor squared, approximately proportional to $1/(1 + Q^2/0.71)^4$. (See Fig. 1.) This factor is responsible for an additional Q^2 -dependence in the case of quasi-elastic scattering and is not included in a ξ -scaling analysis. Thus νW_2^A should show a pronounced scaling violation in the quasi-elastic region.

In the limit of extremely large Q^2 (once the momentum transfer is much larger than the nucleon mass) the variable y converges toward $(-\xi + 1)m$. In this case y and ξ would become equivalent. Significantly, the difference in the elementary cross sections does remain at the q 's of interest and should continue to forbid simultaneous scaling in terms of y and ξ .

Relation between ξ and y In order to better understand the difference between y and ξ scaling, we first discuss the relation between the two variables. Scaling in terms of ξ involves a study of the nuclear response at $\xi=\text{const}$, as a function of the momentum transfer. Fig. 1 shows the relation between ξ and y and it can be seen

that for a given value of ξ , y is a pronounced function of Q^2 , particularly at the lower momentum transfers.




FIG. 1: Relation between ξ and y as a function of momentum transfer Q^2 . The dotted curves (right hand scale) represent the Q^2 -dependence of the Mott cross section (upper curve) and the nucleon form factor squared.

As a consequence, the nuclear DIS scaling function, νW_2^A , at constant value of ξ , includes contributions from a large range of y values when studied over Q^2 . The nuclear quasi-elastic scaling function $F(y)$ (roughly speaking the probability to find in the nucleus nucleons with momentum component y parallel to q) depends strongly on y , as nuclear momentum distributions are a steeply falling function with increasing $|y|$. How ξ -scaling appears in the quasi-elastic region requires an understanding of the behavior of several rapidly varying functions – $\sigma_{eN}(Q^2)$, $\sigma_{Mott}(Q^2)$ and $F(y)$.

III. MODEL STUDIES

In order to understand expose the behavior of the different scaling variables and scaling functions, we made a study of the nuclear response using a simple model. In this model, we describe the inclusive nuclear response as a sum of quasi-elastic scattering, excitation of the nucleon resonances and DIS on the nucleon.

For quasi-elastic scattering our model is based on y -scaling at intermediate momentum transfers. $F(y)$ is determined from the experimental data through

$$F(y) = \frac{d^2\sigma^{\text{exp}}}{d\Omega dE'} / \sigma_{ei} \cdot K$$

where σ_{ei} is the appropriate combination of neutron and proton elastic cross sections and $K = |\mathbf{q}| / \sqrt{m^2 + (y + |\mathbf{q}|)^2}$ is the Jacobian resulting from the change of variables. We fit $F(y)$ with the sum of a Gaussian and an exponential function and limit the fit essentially to the $y < 0$ region. As long as we can neglect the effect of the nucleon removal energy E (beyond the

average $\bar{E}$ which we do take into account) the response is symmetric about $y = 0$. The resulting momentum distribution, $n(k) = -\frac{1}{2\pi y} \frac{dF(y)}{dy}$ is a good approximation of what is expected from theoretical calculations of $n(k)$.

For the resonance and DIS regions the neutron and proton responses have been parameterized by Bodek and Ritchie [17] by fitting inclusive scattering data on the nucleon and removing, for the case of the neutron, the Fermi motion effects occurring in the deuteron structure function. To calculate the nuclear response, we fold the nucleonic response of Ref. [17] with the nuclear momentum distribution $n(k)$ determined above.

The sum of these contribution reproduces the experimental nuclear responses quite well. Typical deviations between model and data are of the order of 20%.

In what follows below we determine νW_2^A from the inclusive cross section σ (both model and experimental) via

$$\nu W_2^A(Q^2) = \nu \cdot \frac{\sigma}{\sigma_M} \left[1 + 2 \tan^2(\theta/2) \cdot \left(\frac{1 + \nu^2/Q^2}{1 + R} \right) \right]^{-1}$$

where σ_M is the cross section for point like constituents and the ratio of longitudinal to transverse cross section, $R = \sigma_L/\sigma_T = (1 + \nu^2/Q^2)W_2/W_1 - 1$. We take $R = \frac{0.32}{Q^2}$. The model cross sections are the sum of the quasi-elastic and inelastic contributions as described above. We display in Fig. 2 the model scaling function νW_2 for iron (per nucleon) for six different ξ values as a function of the momentum transfer along with data from Jefferson Lab E89-008 [16] and SLAC NE3 [18, 19]. At very large Q^2 $\xi = 1$ corresponds to the top of the quasi-elastic peak ($y = 0, x = 1$) and at lower momentum transfers $\xi = 1$ maps to increasingly more negative values of y (see Fig. 1).

From Fig. 2 we see that the sum of the quasi-elastic response and the combined resonance and DIS contributions scales better than the combined resonance and DIS piece alone. The latter falls steeply at low Q^2 . This fall-off is, to a large degree, just compensated by the quasi-elastic contribution. The reduced Q^2 -dependence of the sum leads to the approximate ξ -scaling observed experimentally.

At fixed ξ , increasing Q^2 shifts y to less negative values, where $F(y)$ is larger. At the same time, this increase in the $F(y)$ is largely offset (for quasielastic scattering) by the nucleon elastic form factors which fall with increasing Q^2 . The DIS contribution to νW_2^A at fixed ξ grows with $F(y)$ ($n(k)$) as Q^2 increases but does not suffer from the Q^2 dependence of the elastic form factors. Taken together, these two incongruous behaviors lead to an approximately constant νW_2^A .

While the shift of y with Q^2 is of purely kinematical origin, the decrease of $F(y)$ with larger $|y|$ (the decrease of $n(k)$ with larger k) depends entirely on the behavior of the nuclear momentum distribution. The magnitude of this fall-off is specific to the nucleus under consideration, although for different nuclei the tails of nuclear momentum distributions at large k are not too different given




FIG. 2: νW_2 at fixed values of ξ as a function of Q^2 for iron. Data from SLAC NE3 and Jefferson Lab E89-008 are plotted along with the results of the model described in the text. Dotted lines reflect the quasi-elastic piece, dashed lines the inelastic and the solid line the sum. The data sets correspond to $\xi = 0.6, 0.75, 0.9, 1.05, 1.2$ and 1.35 .

the identical origin of these tails, the short-range nucleon-nucleon correlations. There is, however, no connection we can discern that relates the physics responsible for the fall-off of $n(k)$ to the Q^2 -dependence of the nucleon form factors and the kinematics that distinguishes ξ from y . From this observation we conclude that the apparent ξ -scaling is simply due to how the quasi-elastic piece and resonance and DIS pieces sum together at fixed ξ and is largely accidental in character.

This point on the dependence on $F(y)$ ($n(k)$) is further emphasized by Fig. 3 which shows our model νW_2 for the deuteron along with data from [16, 20, 21, 22]. In the corresponding range of y or k ($0.3 < k < 0$ GeV/c) the deuteron momentum distribution falls off more quickly with increasing y (or k) than in heavier nuclei. Accordingly, the approach to ξ -scaling is different: the rapid fall-off of $n(k)$ actually better compensates the fall-off of the nucleon form factor with Q^2 , and leads to significantly better scaling in terms of ξ .

IV. CONCLUSIONS

We have studied the experimental observation of ξ -scaling in electron-nucleus scattering in the region of q, ν where the cross section would not be expected to scale as it is dominated quasi-elastic electron-nucleon scattering. We find that the apparent ξ -scaling comes about because the quasi-elastic contribution approximately com-




FIG. 3: νW_2 for the deuteron at fixed values of ξ as a function of Q^2 . Data from SLAC experiments E101, E133, and NE4 and Jefferson Lab E89-008 are plotted along with the results of the model described in the text. Dotted lines reflect the quasi-elastic piece, dashed lines the inelastic and the solid line the sum.

pensates for the failure of resonance and DIS contribution at large ξ and low Q^2 to scale. The two dominant, rapidly varying functions distinguishing the behavior of the nuclear scaling function, νW_2 , as a function of y and ξ are the nuclear momentum distribution at low k and the nucleon form factors at large Q^2 . As these have no obvious physical connection, we conclude that the appearance of an approximate ξ -scaling at large ξ comes about accidentally—the two dominant contributions to the inclusive cross section behave such that their sum shows a Q^2 independence characteristic of scaling, but separately they do not.

It remains to be seen whether the concept of duality — which refers to an *average* over many resonances rather than a single one (the nucleon ground state) — can further elucidate this accidental ξ -scaling.

Acknowledgments

This work was supported by the U.S. Department of Energy and the Schweizerische Nationalfonds.

-
- [1] I. Sick *et al.*, Phys. Rev. Lett., 45:871, 1980.
 - [2] J.D. Bjorken and E.A. Paschos, Phys. Rev., 185:1975, 1969.
 - [3] P.E. Sokol *et al.*, Momentum Distributions, Plenum Press, Edts R. Silver, P.E. Sokol
 - [4] V.F. Sears, Physical Review B, 30:44, 1984.
 - [5] P. Eisenberger and P.M. Platzman, Phys. Rev. A, 2:415, 1970.
 - [6] D. Day *et al.*, Ann. Rev. Nucl. Part. Sci., 40:357, 1990.
 - [7] O. Nachtmann, Nucl. Phys. B, 63:237, 1973.
 - [8] R. K. Ellis *et al.*, QCD and Collider Physics, Cambridge Univ. Press, 1996.
 - [9] S.J. Brodsky *et al.*, Phys. Rev. D, 65:114025, 2002.
 - [10] I. Niculescu *et al.*, Phys. Rev. Lett., 85:1182, 2000.
 - [11] I. Niculescu *et al.*, Phys. Rev. Lett., 85:1186, 2000.
 - [12] E.D. Bloom and F.J. Gilman, Phys. Rev. D, 4:2901, 1971.
 - [13] B.W. Filippone *et al.*, Phys. Rev. C, 45:1582, 1992.
 - [14] O. Benhar and S. Liuti, Phys. Lett. B, 358:173, 1995.
 - [15] O. Benhar *et al.*, Phys. Rev. C, 44:2328, 1991.
 - [16] J. Arrington *et al.*, Phys. Rev. Lett., 82:2056, 1999.
 - [17] A. Bodek and J.L. Ritchie, Phys. Rev. D, 23:1070, 1981.
 - [18] D. Day *et al.*, Phys. Rev. Lett., 59:427, 1987.
 - [19] D. Day *et al.*, Phys. Rev. C, 48:1849, 1993.
 - [20] W.P. Schuetz *et al.*, Phys. Rev. Lett., 38:259, 1977.
 - [21] R.G. Arnold *et al.*, Phys. Rev. Lett., 61:806, 1988.
 - [22] S. Rock *et al.*, Phys. Rev. D, 46:24, 1992.